

## Section 12.10: Taylor & Maclaurin Sries

### Goal

We will introduce Taylor and Maclaurin Series, which are special types of power series.

### Initial discussion

**Recall 1.** A *power series* is an infinite power series of the form

$$\sum_{n=0}^{\infty} c_n(x-a)^n = c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots$$

We say that the power series is centered at  $x = a$ .

**Question 2.** What functions have power series representations?

**Example 3.** Consider the function

$$f(x) = \frac{1}{1-x}.$$

Have we seen this expression in the context of series? Yes, recall that when  $|r| < 1$ , we have the following for geometric series

$$\sum_{n=0}^{\infty} ar^n = \sum_{n=1}^{\infty} ar^{n-1} = \frac{a}{1-r}.$$

This implies that

$$f(x) = \frac{1}{1-x} = \sum_{n=0}^{\infty} x^n,$$

as long as  $|x| < 1$ . That is,  $f$  has a power series representation centered at 0 with radius 1. Outside the interval  $(-1, 1)$  the two expressions are not equal.

Let's take a look at what is going on graphically by looking at the [Power Series and Interval of Convergence Applet](http://calculusapplets.com/), which is available at <http://calculusapplets.com/>.

Not every power series takes the form of a geometric series. So, we need a more general method.

### Taylor & Maclaurin Series

Suppose  $f$  has a power series representation (with  $|x-a| < R$ ):

$$f(x) = \sum_{n=0}^{\infty} c_n(x-a)^n = c_0 + c_1(x-a) + c_2(x-a)^2 + \cdots$$

Since these functions are equal, their derivatives agree:

$$f'(x) = \sum_{n=1}^{\infty} n c_n(x-a)^{n-1} = c_1 + 2c_2(x-a) + 3c_3(x-a)^2 + \cdots$$

This implies that

$$f'(a) = c_1 + 0 + 0 + \cdots = c_1.$$

That is,

$$c_1 = \underline{\hspace{2cm}}$$

Let's repeat this process with the second derivative. We see that

$$f''(x) = \sum_{n=2}^{\infty} n(n-1)c_n(x-a)^{n-2} = \underline{\hspace{15cm}}$$

This implies that

$$f''(a) = \underline{\hspace{4cm}} = \underline{\hspace{4cm}}.$$

Then

$$c_2 = \underline{\hspace{2cm}}$$

And again using the third derivative:

$$f'''(x) = \sum_{n=3}^{\infty} n(n-1)(n-2)c_n(x-a)^{n-3} = \underline{\hspace{15cm}}$$

This implies that

$$f'''(a) = \underline{\hspace{4cm}} = \underline{\hspace{4cm}}.$$

Then

$$c_3 = \underline{\hspace{2cm}}$$

If we continue this way, we'll obtain the following

$$c_n = \underline{\hspace{2cm}}$$

**Note 4.** Here are a couple of conventions:

1.  $0! = 1! = 1$
2.  $f^{(0)}(a) = f(a)$

**Theorem 5.** *If  $f$  has a power series representation at  $x = a$ , then it must be of the form*

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for  $|x - a| < R$ , where  $R$  is the radius of convergence.

The above power series is called the *Taylor Series of  $f$  centered at  $x = a$* . In the special case that  $a = 0$ , we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n,$$

which is called the *Maclaurin series of  $f$* .

**Important Note 6.** We can always compute a Taylor/Maclaurin series, but that does not mean that it is equal to the given function. We only know that a Taylor/Maclaurin Series is equal to a given function *if* the given function can be represented by a power series. We will only deal with these types of functions.

**Example 7.**

- (a) Find Maclaurin series for  $f(x) = e^x$  and its radius of convergence (given that  $f$  has a power series representation).

- (b) Find Maclaurin series for  $f(x) = \sin x$  and its radius of convergence (given that  $f$  has a power series representation).

## Common Taylor Series

Here are some common Taylor Series.

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n \quad R = 1$$

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad R = \infty$$

$$\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} \quad R = \infty$$

$$\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \quad R = \infty$$

$$\arctan x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1} \quad R = 1$$

## Taylor Series approximation

If a function has a Taylor Series representation, then we can use a finite number of terms to approximate the function. We define the *k*th degree Taylor polynomial of *f* at  $x = a$  to be

$$T_k(x) = \sum_{n=0}^k \frac{f^{(n)}(a)}{n!} (x - a)^n$$

Let's take a look at the [Taylor Series and Polynomials Applet](http://calculusapplets.com/) available at <http://calculusapplets.com/>.

### Example 8.

(a) Use the 9th degree Taylor polynomial for  $\arctan x$  to approximate  $\pi$ .

(b) Approximate the following integral using a 5th degree Taylor polynomial for  $\sin x$ .

$$\int_0^1 x \sin(x^3) dx$$