

## 3 Relations and Functions

### 3.1 Relations

**Definition 3.1.** An **ordered pair** is an object of the form  $(x, y)$ . Two ordered pairs  $(x, y)$  and  $(a, b)$  are **equal** if  $x = a$  and  $y = b$ .

**Definition 3.2.** An  **$n$ -tuple** is object of the form  $(x_1, x_2, \dots, x_n)$ . Each  $x_i$  is referred to as the  $i$ th **component**.

Note that an ordered pair is just a 2-tuple.

**Definition 3.3.** If  $X$  and  $Y$  are sets, the **Cartesian product** of  $X$  and  $Y$  is defined by

$$X \times Y = \{(x, y) : x \in X, y \in Y\}.$$

That is,  $X \times Y$  is the set of all ordered pairs where the first element is from  $X$  and the second element is from  $Y$ . The set  $X \times X$  is sometimes denoted by  $X^2$ . We similarly define the Cartesian product of  $n$  sets, say  $X_1, \dots, X_n$ , by

$$\prod_{i=1}^n X_i = X_1 \times \dots \times X_n = \{(x_1, \dots, x_n) : \text{each } x_i \in X_i\}.$$

**Example 3.4.** Let  $A = \{a, b, c\}$  and  $B = \{\ominus, \odot\}$ . Then

$$A \times B = \{(a, \ominus), (a, \odot), (b, \ominus), (b, \odot), (c, \ominus), (c, \odot)\}.$$

**Exercise 3.5.** Using the sets  $A$  and  $B$  from the previous example, find  $B \times A$ .

**Exercise 3.6.** Using the set  $B$  from the previous examples, find  $B \times B$ .

**Exercise 3.7.** What general conclusion can you make about  $X \times Y$  versus  $Y \times X$ ? When will they be equal?

**Exercise 3.8.** If  $X$  and  $Y$  are both finite sets, then how many elements will  $X \times Y$  have? Be as specific as possible.

**Exercise 3.9.** Let  $A = \{1, 2, 3\}$ ,  $B = \{1, 2\}$ , and  $C = \{1, 3\}$ . List the elements of the set  $A \times B \times C$ .

**Exercise 3.10.** Let  $A = \mathbb{N}$  and  $B = \mathbb{R}$ . Describe the elements of the set  $A \times B$ .

**Exercise 3.11.** Let  $A$  be the set of all differentiable functions on the open interval  $(0, 1)$ , and let  $B$  equal the set of all derivatives of functions in  $A$  evaluated at  $x = \frac{1}{2}$ . Describe the elements of the set  $A \times B$ .

**Exercise 3.12.** Three space,  $\mathbb{R}^3$ , is a Cartesian product. Unpack the meaning of  $\mathbb{R}^3$  using the Cartesian product, and write the complete set notation version.

**Exercise 3.13.** Let  $X = [0, 1]$  and let  $Y = \{1\}$ . Describe geometrically what  $X \times Y$ ,  $Y \times X$ ,  $X \times X$ , and  $Y \times Y$  look like.

**Definition 3.14.** Let  $X$  and  $Y$  be sets. A **relation** from a set  $X$  to a set  $Y$  is a subset of  $X \times Y$ . A relation on  $X$  is a subset of  $X \times X$ .

**Example 3.15.** You may not realize it, but you are familiar with many relations. For example, on the real numbers, we have the relation  $\leq$ . We could say that  $(3, \pi)$  is in the relation since  $3 \leq \pi$ . However,  $(1, -1)$  is not in the relation since  $1 \not\leq -1$ . (Order matters!)

**Remark 3.16.** Different notations for relations are used in different contexts. When talking about relations in the abstract, we indicate that a pair  $(a, b)$  is in the relation by some notation like  $a \sim b$ , which is read “ $a$  is related to  $b$ .”

**Example 3.17.** Let  $P_f$  denote the set of all people with accounts on Facebook. Define  $F$  via  $xFy$  iff  $x$  is friends with  $y$ . Then  $F$  is a relation on  $P_f$ .

**Remark 3.18.** We can often represent relations using graphs or digraphs. Given a finite set  $X$  and a relation  $\sim$  on  $X$ , a **digraph** (short for *directed graph*) is a discrete graph having the members of  $X$  as vertices and a directed edge from  $x$  to  $y$  iff  $x \sim y$ .

**Example 3.19.** When we write  $x^2 + y^2 = 1$ , we are implicitly defining a relation. In particular, the relation is the set of ordered pairs  $(x, y)$  satisfying  $x^2 + y^2 = 1$ . In set notation:

$$\{(x, y) : x^2 + y^2 = 1\}$$

The graph of this relation in  $\mathbb{R}^2$  is the standard unit circle.

**Exercise 3.20.** Define  $\sim$  on  $\mathbb{R}^2$  via  $x \sim y$  iff  $x \leq y$ . Draw a picture of this relation in  $\mathbb{R}^2$ .

**Example 3.21.** Let  $A = \{a, b, c\}$  and define  $\sim = \{(a, a), (a, b), (b, c), (c, b), (c, a)\}$ . The digraph for  $\sim$  is a graph with vertices  $a, b, c$  and the following arrows:  $a$  to  $a$ ,  $a$  to  $b$ ,  $b$  to  $c$ ,  $c$  to  $b$ ,  $c$  to  $a$ .

**Exercise 3.22.** Let  $A = \{1, 2, 3, 4, 5, 6\}$  Define  $|$  on  $A$  via  $x|y$  iff  $x$  divides  $y$ . Draw the digraph for  $|$  on  $A$ .

**Definition 3.23.** Let  $\sim$  be a relation on a set  $A$ .

1.  $\sim$  is **reflexive** if for all  $x \in A$ ,  $x \sim x$  (every element is related to itself).
2.  $\sim$  is **symmetric** if for all  $x, y \in A$ , if  $x \sim y$ , then  $y \sim x$ .
3.  $\sim$  is **transitive** if for all  $x, y, z \in A$ , if  $x \sim y$  and  $y \sim z$ , then  $x \sim z$ .

**Example 3.24.**

1.  $\leq$  on  $\mathbb{R}$  is reflexive and transitive, but not symmetric.  $<$  on  $\mathbb{R}$  is transitive, but not symmetric and not reflexive.
2. If  $S$  is a set, then  $\subseteq$  on  $\mathcal{P}(S)$  is reflexive and transitive, but not symmetric.
3.  $=$  on  $\mathbb{R}$  is reflexive, symmetric, and transitive.

**Exercise 3.25.** Given a finite set  $A$  and a relation  $\sim$ , describe what each of reflexive, symmetric, and transitive look like in terms of a digraph.

**Exercise 3.26.** Let  $P$  be the set of people at a party and define  $N$  via  $(x, y) \in N$  iff  $x$  knows the name of  $y$ . Describe what it would mean for  $N$  to be reflexive, symmetric, and transitive.

**Exercise 3.27.** Determine whether each of the following relations are reflexive, symmetric, or transitive.

1. Let  $P_f$  denote the set of all people with accounts on Facebook. Define  $F$  via  $xFy$  iff  $x$  is friends with  $y$ .
2. Let  $P$  be the set of all people and define  $H$  via  $xHy$  iff  $x$  and  $y$  have the same height.
3. Let  $P$  be the set of all people and define  $T$  via  $xTy$  iff  $x$  is taller than  $y$ .
4. Consider the relation “divides” on  $\mathbb{Z}$ .
5. Let  $L$  be the set of lines and define  $\parallel$  via  $l_1 \parallel l_2$  iff  $l_1$  is parallel to  $l_2$ .
6. Let  $C[0, 1]$  be the set of continuous functions on  $[0, 1]$ . Define  $f \sim g$  iff

$$\int_0^1 |f(x)| dx = \int_0^1 |g(x)| dx.$$

7. Define  $\sim$  on  $\mathbb{N}$  via  $n \sim m$  iff  $n + m$  is even.
8. Define  $D$  on  $\mathbb{R}$  via  $(x, y) \in D$  iff  $x = 2y$ .