

The Temperley-Lieb Algebras of Types A and B and Their Associated Diagram Algebras

Dana Ernst

University of Colorado, Boulder
Department of Mathematics

Slow Pitch: October 10, 2007

Definition

Let n be a positive integer. The *Temperley-Lieb Algebra of Type A*, $\mathrm{TL}_n(A)$, with parameter δ is defined to be the associative, unital algebra over the ring $\mathbb{Z}[\delta]$ generated by elements e_1, e_2, \dots, e_{n-1} subject only to the relations

$$e_i^2 = \delta e_i, \text{ for all } i$$

$$e_i e_j = e_j e_i, \text{ for } |i - j| \geq 2$$

$$e_i e_j e_i = e_i, \text{ for } |i - j| = 1$$

Theorem

$\mathrm{TL}_n(A)$ is a finite dimensional associative algebra over $\mathbb{Z}[\delta]$. A basis may be described in terms of “reduced words” in the algebra generators e_j . The rank of $\mathrm{TL}_n(A)$ is the n th Catalan number:


$$C_n = \frac{1}{n+1} \binom{2n}{n} = \frac{(2n)!}{(n+1)!n!}.$$

Some Remarks:

- ▶ $TL_n(A)$ was invented in 1971 by Temperley and Lieb.
- ▶ First arose in the context of integrable Potts models in statistical mechanics.
- ▶ As well as having applications in physics, $TL_n(A)$ appears in the framework of knot theory, braid groups, Coxeter groups and their corresponding Hecke algebras, and subfactors of von Neumann algebras.
- ▶ Penrose/Kauffman use diagram algebra to model $TL_n(A)$ in 1971.
- ▶ In 1987, Vaughan Jones recognized that $TL_n(A)$ is isomorphic to a particular quotient of the Hecke Algebra of type A_{n-1} (the symmetric group, S_n).

Definition

A *standard n -box* is a rectangle with $2n$ nodes, labeled as follows:




An *n -diagram* is a graph drawn on the nodes of a standard n -box such that


- ▶ Every node is connected to exactly one other node by a single edge.
- ▶ All edges must be drawn inside the n -box.
- ▶ The graph can be drawn so that no edges cross.

Example

Here is an example of a 5-diagram.




Here is another.



Example

Here is an example that is *not* a diagram.




Definition

The associative *diagram algebra*, $\mathcal{D}_n(A)$, is the free $\mathbb{Z}[\delta]$ -module having the set of n -diagrams as a basis with multiplication defined as follows.

If d and d' are n -diagrams, then dd' is obtained by identifying the “south face” of d with the “north face” of d' , and then replacing any closed loops with a factor of δ .


Example

Multiplication of two 5-diagrams.




Example

Here's another example.




Example

And here's one more.



$$= \delta^3$$




Theorem


The rank of the diagram algebra $\mathcal{D}_n(A)$ is C_n .

Proof.


The number of sequences of n pairs of well-formed parentheses is C_n . There is a one-to-one correspondence between n -diagrams and sequences of n pairs of well-formed parentheses.



Now, we define a few “simple” n -diagrams. Let



\vdots



\vdots



Claim 1: The diagrams d_1, d_2, \dots, d_{n-1} generate $\mathcal{D}_n(A)$.

Claim 2: The generators d_1, d_2, \dots, d_{n-1} satisfy the relations of $\text{TL}_n(A)$.


For all i , we have

$$\begin{aligned} d_i^2 &= \begin{array}{c} \begin{array}{cccccccc} 1 & \dots & i-1 & i & i+1 & i+2 & \dots & n \end{array} \\ \begin{array}{|c|} \hline \begin{array}{cccccccc} \bullet & & \bullet & \text{---} & \bullet & \bullet & \bullet & \bullet \end{array} \\ \hline \begin{array}{cccccccc} \bullet & & \bullet & \text{---} & \bullet & \bullet & \bullet & \bullet \end{array} \\ \hline \begin{array}{cccccccc} \bullet & & \bullet & \text{---} & \bullet & \bullet & \bullet & \bullet \end{array} \\ \hline \end{array} \\ \dots \\ &= \delta \begin{array}{c} \begin{array}{cccccccc} 1 & \dots & i-1 & i & i+1 & i+2 & \dots & n \end{array} \\ \begin{array}{|c|} \hline \begin{array}{cccccccc} \bullet & & \bullet & \text{---} & \bullet & \bullet & \bullet & \bullet \end{array} \\ \hline \begin{array}{cccccccc} \bullet & & \bullet & \text{---} & \bullet & \bullet & \bullet & \bullet \end{array} \\ \hline \end{array} \\ \dots \\ &= \delta d_i \end{aligned}$$

For $|i - j| \geq 2$, we have

$$\begin{aligned}
 d_i d_j &= \begin{array}{c} \begin{array}{cccccccccccccccc} 1 & \dots & i-1 & i & i+1 & i+2 & \dots & j-1 & j & j+1 & j+2 & \dots & n \end{array} \\ \left[\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right] \\ \begin{array}{cccccccccccccccc} \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots \\ \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots \\ \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots \end{array} \end{array} \\
 &= \begin{array}{c} \begin{array}{cccccccccccccccc} 1 & \dots & i-1 & i & i+1 & i+2 & \dots & j-1 & j & j+1 & j+2 & \dots & n \end{array} \\ \left[\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \right] \\ \begin{array}{cccccccccccccccc} \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots \\ \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots \\ \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots & \text{---} & \text{---} & \vdots & & \vdots \end{array} \end{array} \\
 &= d_j d_i
 \end{aligned}$$

For $|i - j| = 1$ (here, $j = i + 1$; $j = i - 1$ being similar), we have



Claim 1 and Claim 2, along with the fact that $\text{TL}_n(A)$ and $\mathcal{D}_n(A)$ have the same dimension, suggest the following theorem.

Theorem

$\text{TL}_n(A)$ and $\mathcal{D}_n(A)$ are isomorphic as $\mathbb{Z}[\delta]$ -algebras under the correspondence

$$e_j \mapsto d_j.$$

Now, consider the group algebra of the symmetric group S_n over \mathbb{Z} :

$$\mathbb{Z}[S_n]$$

Recall that S_n is generated by the adjacent transpositions:

$$(1\ 2), (2\ 3), \dots, (n-1\ n).$$

Define

$$s_i = (i\ i+1).$$

Next, take the (principal) ideal, J , of $\mathbb{Z}[S_n]$ generated by all elements of the form

$$1 + s_i + s_j + s_i s_j + s_j s_i + s_i s_j s_i,$$

where $|i - j| = 1$ (i.e., s_i and s_j are noncommuting generators).

Definition

Let $\sigma = s_{i_1} \dots s_{i_r} \in S_n$ be reduced. We say that σ is *fully commutative*, or *FC*, if any two reduced expressions for σ may be obtained from each other by repeated commutation of adjacent generators. In other words, σ has no reduced expression containing $s_i s_j s_i$ for $|i - j| = 1$.

Example

$s_1 s_2 s_4 s_1 = (1\ 2)(2\ 3)(4\ 5)(1\ 2)$ is a reduced expression for an element in S_5 . This element is *not* FC.

$$s_1 s_2 s_4 s_1 = s_1 s_2 s_1 s_4$$

Now, let

$$b_{s_i} = (1 + s_i) + J \in \mathbb{Z}[S_n]/J.$$

Theorem

As a unital algebra, $\mathbb{Z}[S_n]/J$ is generated by $b_{s_1}, \dots, b_{s_{n-1}}$.

Definition

If $\sigma = s_{i_1} \dots s_{i_r}$ is reduced and FC, then

$$b_\sigma = b_{s_{i_1}} \dots b_{s_{i_r}}$$

is a well-defined element of $\mathbb{Z}[S_n]/J$.

Theorem

The set $\{b_\sigma : \sigma \text{ FC}\}$ is a free \mathbb{Z} -basis for $\mathbb{Z}[S_n]/J$.

That is, $\mathbb{Z}[S_n]/J$ has a basis indexed by the fully commutative elements of S_n .

If we let $\delta = 2$, we have the following result.

Theorem

The algebras $\mathbb{Z}[S_n]/J$ and $\text{TL}_n(A)$ (with $\delta = 2$) are isomorphic as \mathbb{Z} -algebras under the correspondence

$$b_{s_i} = (1 + s_i) + J \mapsto d_i.$$