

## Section 11.2: Calculus with Parametric Curves

### Goal

In this section, we discuss derivatives of parametric curves and learn how to find area, arc length, and surface area in the context of parametric curves.

### Derivatives of parametric curves

Suppose that

$$x = f(t)$$

$$y = g(t)$$

define a parametric curve. If we could eliminate the parameter, we would end up with something of the form

$$y = F(x)$$

(where  $F$  is *not* an antiderivative, but rather some function of  $x$ ).

If we were to substitute back in, we obtain

$$\frac{dy}{dt} = \frac{dF(f(t))}{dt}$$

By the chain rule (assuming  $F$ ,  $f$ , and  $g$  are differentiable), we get

$$\frac{dy}{dt} = \frac{dF}{dx} \cdot \frac{dx}{dt}$$

Now, as long as  $f'(t) \neq 0$ , we obtain

$$F'(x) = F'(f(t)) = \frac{dy/dt}{dx/dt} = \frac{g'(t)}{f'(t)}$$

That is,

**Theorem 1.**

$$\frac{dy}{dx} = \frac{g'(t)}{f'(t)} \quad (\text{provided } dx/dt \neq 0).$$

**Important Note 2.**

1. This formula allows us to find derivative of  $y$  with respect to  $x$  without actually having to eliminate the parameter.
2. Horizontal tangents when  $dy/dt = 0$  (and  $dx/dt \neq 0$ ).
3. Vertical tangents when  $dx/dt = 0$  (and  $dy/dt \neq 0$ ).

**Example 3.**

- (a) Define the parametric curve  $C : x = t^2, y = t^3 - 3t$ . Find points where  $C$  has (i) horizontal tangents, and (ii) vertical tangents.

- (b) Define the parametric curve  $C : x = 2 \sin 2t, y = 3 \sin t$ . Find slope of both tangent lines at the point  $(0, 0)$ .

**Note 4.** To find second derivative:

$$\frac{d^2y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \underline{\hspace{2cm}} \neq \frac{d^2y/dt^2}{d^2x/dt^2}.$$

## Area, arc length, and surface area

**Recall 5.**

- Area:  $A = \int_a^b y \, dx$
- Arc length:  $s = \int_a^b \sqrt{1 + [dy/dx]^2} \, dx$
- Surface area:  $S = 2\pi \int_a^b r(x) \sqrt{1 + [dy/dx]^2} \, dx$

If a parametric curve  $C$  is given by

$$x = f(t)$$

$$y = g(t)$$

then

$dx = \underline{\hspace{2cm}}$ $dy = \underline{\hspace{2cm}}$
---

and

$$\begin{aligned} \sqrt{1 + [dy/dx]^2} \, dx &= \sqrt{1 + \left( \frac{dy/dt}{dx/dt} \right)^2} f'(t) dt \\ &= \sqrt{[f'(t)]^2 + [g'(t)]^2} \, dt. \end{aligned}$$

That is,

$\sqrt{1 + [dy/dx]^2} \, dx = \sqrt{[dx/dt]^2 + [dy/dt]^2} \, dt$
---

If  $C$  is *smooth* (i.e.,  $f'$  and  $g'$  are continuous), then by making the appropriate substitutions, we can find area, arc length, and surface area in the context of parametric curves.

**Important Note 6.**

1. If  $C$  is on the interval  $[\alpha, \beta]$ , then for *area*, we integrate from either  $t = \alpha \rightarrow t = \beta$  or  $t = \beta \rightarrow t = \alpha$ ; the proper choice being the one that corresponds to traversing curve from  $L \rightarrow R$ .
2. For all integrals, we need to make sure we trace out curve *exactly once* from  $\alpha \rightarrow \beta$  (otherwise, we may get too much or too little).

**Example 7.**

(a) Find area under  $C : x = e^{3t}, y = e^{-t}$  on  $[0, \ln 2]$ .

(b) Find arc length of  $C : x = \cos t, y = \sin t$  on  $[0, 2\pi]$ .

(c) Find surface area of unit sphere.